

LICE TREATMENT OPTIONS					
Product and Cost	Prescription or Over the counter	Active Ingredient	Kills live lice and unhatched eggs	Combing required	Retreatment required
Lice MD \$9.98	Over-the-counter	Non-Pesticide-Silicone based Dimethicone	Kills live lice not unhatched eggs	Combing required. Any missed eggs or live lice can lead to recurrence.	Inspect hair daily for 10 days and repeat as needed.
Lice of Clinics Of America--At home treatment \$35.00 (FREE head lice checks)	Over-the-counter Shampoo and comb sold at Lice Clinics of America 970.233.8787	Non-Pesticide-Silicone based Dimethicone	Kills live lice not Unhatched eggs	Combing required. Any missed eggs or live lice can lead to recurrence.	Inspect hair daily For 10 days and repeat as needed.
Rid \$9.97	Over the counter	Pesticide-Pyrethrin	Kills live lice not unhatched eggs	Combing required. Any missed eggs or live lice can lead to recurrence.	Second treatment is needed in 9 days to kill any newly hatched lice.
Nix \$10.44	Over the counter	Pesticide-Synthetic pyrethrin	Kills live lice not unhatched eggs	Combing required. Any missed eggs or live lice can lead to recurrence.	Second treatment is needed in 9 days to kill any newly hatched lice.
Ulesfia lotion \$210.00 (may be covered by insurance)	By prescription only	Non-pesticide-Benzyl alcohol lotion	Kills live lice not unhatched eggs	Combing required. Any missed eggs or live lice can lead to recurrence.	Second treatment is needed in 7 days to kill any newly hatched lice.
Sklice \$320.00 (may be covered by insurance)	By prescription only	Pesticide-Ivermectin lotion	Prevents newly hatched lice from surviving	1 combing required	Effective in most patients with a single application.
Ovide \$95.00 (may be covered by insurance)	By prescription only	Pesticide-Malathion	Kills some live lice and some lice eggs	Combing required. Any missed eggs or live lice can lead to recurrence.	Second treatment recommended if lice are present
Lice Clinics of America \$195 (Medicaid Discount \$99)	By appointment 970.233.8787 Provide free lice checks.	Hot air device	Kills live lice and eggs in a single treatment	None	None

- Do not use extra amounts of the medication unless instructed to do so by your physician and pharmacist. Insecticide products can be dangerous if they are misused or overused.
- All medications/products should be kept out of the eyes. If it gets into the eyes, it should be immediately flushed.
- Do not treat an infested person more than 2-3 times with the same medication if it does not seem to be working. This may be caused by using the medication incorrectly or by resistance to the medicine/product. Always seek the advice of your health care provider if this should happen. An alternative medication/product may be advised.
- Do not use different lice medication at the same time unless instructed to do so by your physician and pharmacist.